Author Index

Abberton, M.T. *see* Williams, T.A.
Agnusdei, M.G. and Mazzanti, A. Frequency of defoliation of native and naturalized species of the Flooding Pampas (Argentina) 344
Arias, C. *see* Córdoba, A.

Baker, J.E. *see* Callow, M.N.
Banfield, J.D. *see* Lenssen, A.W.
Barrett, P.D., Laidlaw, A.S., Mayne, C.S. and Christie, H. Pattern of herbage intake rate and bite dimensions of rotationally grazed dairy cows as sward height declines 362
Binnie, R.C. *see* Frost, J.P.
Binnie, R.C., Mayne, C.S. and Laidlaw, A.S. The effects of rate and timing of application of fertilizer nitrogen in late summer on herbage mass and chemical composition of perennial ryegrass swards over the winter period in Northern Ireland 46
Bruckental, I. *see* Ungar, E.D.

Callow, M.N., Baker, J.E. and Hough, G.M. The effect of accumulation period and harvest date in spring on dry-matter yield and forage quality in mixed swards containing *Lolium* spp. and *Trifolium subterraneum* in Western Australia 10
Carrère, P. *see* Parsons, A.J.
Carrère, P., Louault, F., De Faccio Carvalho, P.C., Lafarge, M. and Soussana, J.F. How does the vertical and horizontal structure of a perennial ryegrass and white clover sward influence grazing? 118
Cash, S.D. *see* Lenssen, A.W.
Chen, Z.Z. *see* Wang, S.P.
Christie, H. *see* Barrett, P.D.
Cooper, A. *see* Weller, R.F.
Córdoba, A., García Seffino, L., Moreno, H., Arias, C., Grunberg, K., Zenoff, A. and Taleisnik, E. Characterization of the effect of high salinity on roots of *Chloris gayana* Kunth: carbohydrate and lipid accumulation and growth 162

Davies, D.A. *see* Fothergill, M.
Davies, D.R. *see* Fraser, M.D.
Davies, D.R. *see* Miller, L.A.
De Faccio Carvalho, P.C. *see* Carrère, P.
Delpy, R. *see* Loiseau, P.
Dewhurst, R.J., Scollan, N.D., Youell, S.J., Tweed, J.K.S. and Humphreys, M.O. Influence of species, cutting date and cutting interval on the fatty acid composition of grasses 68
Driehuis, F. and van Wikselaar, P.G. Determination of microbial protein in perennial ryegrass silage 75
Driehuis, F., Oude Elferink, S.J.W.H. and Van Wikselaar, P.G. Fermentation characteristics and aerobic stability of grass silage inoculated with *Lactobacillus buchneri*, with or without homofermentative lactic acid bacteria 330

Eriksen, J. Implications of grazing by sows for nitrate leaching from grassland and the succeeding cereal crop 317

Fehmi, J.S., Laca, E.A. and Rice, K.J. The effect of small gaps in California annual grassland on above-ground biomass production 323
Fothergill, M., Davies, D.A. and Morgan, C.T. Extensification of grassland use in the Welsh uplands: sheep performance in years 1–6 105
Fraser, M.D., Fychan, R. and Jones, R. The effect of harvest date and inoculation on the yield, fermentation characteristics and feeding value of forage pea and field bean silages 218
Fraser, M.D., Winters, A., Fychan, R., Davies, D.R. and Jones, R. The effect of harvest date and inoculation on the yield, fermentation characteristics and feeding value of kale silage 151
Frost, J.P. and Binnie, R.C. Effect of herbage mass per unit area and conditioning treatment on herbage drying rate 169
Fychan, R. *see* Fraser, M.D.
Fychan, R. *see* Winters, A.L.

García Seffino, L. *see* Córdoba, A.
Gilliland, T.J. and Mann, R.L. The silage yield potential of hybrid ryegrass over a 5-year period in Northern Ireland relative to other ryegrasses and timothy 305
Grunberg, K. *see* Córdoba, A.

Hameleers, A. *see* Leach, K.A.
Haneklaus, S. *see* Wang, S.P.
Harvey, B.M.R. *see* Wilson, F.E.A.
Haugland, E. and Tawfiq, M. Root and shoot competition between established grass species and newly sown seedlings during spring growth 193

Herrmann, A. and Schachtel, G.A. OSYAQ, an organ-specific growth model for forage grasses 268
Hervás, G. *see* Mould, F.L.
Hough, G.M. *see* Callow, M.N.
Humphreys, M.O. *see* Dewhurst, R.J.
Humphreys, M.O. *see* Miller, L.A.

Iglesias, I. *see* Lloveras, J.

Järvenranta, K. *see* Virkajärvi, P.
Jauhiainen, L. *see* Niemeläinen, O.
Jefferson, P.G., Lawrence, T. and Kielly, G.A. Divergent recurrent selection for seedling tiller number in Altai wildrye 285
Jones, J.R. *see* Wright, I.A.
Jones, R. *see* Fraser, M.D.
Jones, R. *see* Winters, A.L.

Kielly, G.A. *see* Jefferson, P.G.

Laca, E.A. *see* Fehmi, J.S.
Lafarge, M. *see* Carrère, P.
Laidlaw, A.S. *see* Barrett, P.D.
Laidlaw, A.S. *see* Binnie, R.C.
Lawrence, T. *see* Jefferson, P.G.
Leach, K.A. and Hameleers, A. The effects of a foliar spray containing phosphorus and zinc on the development, composition and yield of forage maize (Research Note) 311
Leaver, J.D. *see* Pulido, R.G.
Lenssen, A.W., Banfield, J.D. and Cash, S.D. The influence of trichome density on the drying rate of alfalfa forage 1
Lloveras, J. and Iglesias, I. Morphological development and forage quality changes in crimson clover (*Trifolium incarnatum* L.) 395
Loiseau, P., Soussana, J.-F., Louault, F. and Delpy, R. Soil N contributes to the oscillations of the white clover content in mixed swards of perennial ryegrass under conditions that simulate grazing over five years 205
Louault, F. *see* Carrère, P.
Louault, F. *see* Loiseau, P.

MacRae, J.C. *see* Miller, L.A.
Mann, R.L. *see* Gilliland, T.J.
Marrs, R.H. *see* Petrov, P.
Mayne, C.S. *see* Barrett, P.D.
Mayne, C.S. *see* Binnie, R.C.
Mayne, C.S. *see* Sayers, H.J.
Mazzanti, A. *see* Agnusdei, M.G.
McAdam, J.H. *see* Wilson, F.E.A.
Miettinen, E. *see* Niemeläinen, O.
Miller, L.A., Moorby, J.M., Davies, D.R., Humphreys, M.O., Scollan, N.D., MacRae, J.C. and Theodorou, M.K. Increased concentration of water-soluble carbohydrate in perennial ryegrass (*Lolium perenne* L.): milk production from late-lactation dairy cows 383
Moorby, J.M. *see* Miller, L.A.
Moreno, H. *see* Córdoba, A.
Morgan, C.T. *see* Fothergill, M.
Mould, F.L., Hervás, G., Owen, E., Wheeler, T.R., Smith, N.O. and Summerfield, R.J. The effect of cultivar on the rate and extent of combining pea straw degradability examined *in vitro* using the Reading Pressure Technique 374
Murray, R.S. *see* Porter, M.G.

Niemeläinen, O., Jauhiainen, L. and Miettinen, E. Yield profile of tall fescue (*Festuca arundinacea*) in comparison with meadow fescue (*F. pratensis*) in Finland 249

Orr, R.J., Rutter, S.M., Penning, P.D. and Rook, A.J. Matching grass supply to grazing patterns for dairy cows 352
Oude Elferink, S.J.W.H. *see* Driehuis, F.
Owen, E. *see* Mould, F.L.

Parsons, A.J. *see* Wright, I.A.
Parsons, A.J., Schwinning, S. and Carrère, P. Plant growth functions and possible spatial and temporal scaling errors in models of herbivory 21
Penning, P.D. *see* Orr, R.J.
Penning, P.D. *see* Yarrow, N.H.
Petrov, P. and Marrs, R.H. The reclamation of bracken-dominated pastures in Bulgaria using asulam and fertilizers 131
Porter, M.G. and Murray, R.S. The volatility of components of grass silage on oven drying and the interrelationship between dry-matter content estimated by different analytical methods 405
Pulido, R.G. and Leaver, J.D. Quantifying the influence of sward height, concentrate level and initial milk yield on the milk production and grazing behaviour of continuously stocked dairy cows 57

Ravid, N. *see* Ungar, E.D.
Rawson, H.M. *see* Turner, N.C.
Rhodes, I. *see* Williams, T.A.
Rice, K.J. *see* Fehmi, J.S.
Rook, A.J. *see* Orr, R.J.
Rutter, S.M. *see* Orr, R.J.

Sayers, H.J. and Mayne, C.S. Effect of early turnout to grass in spring on dairy cow performance 259
Schachtel, G.A. *see* Herrmann, A.

Schnug, E. *see* Wang, S.P.
Schwinning, S. *see* Parsons, A.J.
Scollan, N.D. *see* Dewhurst, R.J.
Scollan, N.D. *see* Miller, L.A.
Smith, N.O. *see* Mould, F.L.
Soussana, J.F. *see* Carrère, P.
Soussana, J.-F. *see* Loiseau, P.
Summerfield, R.J. *see* Mould, F.L.

Taleisnik, E. *see* Córdoba, A.
Tawfiq, M. *see* Haugland, E.
Theodorou, M.K. *see* Miller, L.A.
Thomson, C.J. *see* Turner, N.C.
Thornley, W. *see* Williams, T.A.
Turner, N.C., Thomson, C.J. and Rawson, H.M. Effect of temperature on germination and early growth of subterranean clover, capeweed and Wimmera ryegrass 97
Tweed, J.K.S. *see* Dewhurst, R.J.

Ungar, E.D., Ravid, N. and Bruckental, I. Bite dimensions for cattle grazing herbage at low levels of depletion 35

Virkajärvi, P. and Järvenranta, K. Leaf dynamics of timothy and meadow fescue under Nordic conditions 294

Walton, D.W.H. *see* Wilson, F.E.A.
Wang, S.P., Wang, Y.F., Chen, Z.Z., Schnug, E. and Haneklaus, S. Sulphur concentration of soils and plants and its requirement for ruminants in the Inner Mongolia steppe of China (Research Note) 418
Wang, Y.F. *see* Wang, S.P.
Weller, R.F. and Cooper, A. Seasonal changes in the crude protein concentration of mixed swards of white clover/perennial ryegrass grown without fertilizer N in an organic farming system in the United Kingdom (Research Note) 92
Wheeler, T.R. *see* Mould, F.L.
van Wikselaar, P.G. *see* Driehuis, F.
Williams, T.A., Abberton, M.T., Thornley, W. and Rhodes, I. Relationships between the yield of perennial ryegrass and of small-leaved white clover under cutting or continuous grazing by sheep 231
Williams, T.A., Abberton, M.T., Thornley, W. and Rhodes, I. No relationship between leaf size and yield in medium leaf size white clover varieties under rotational sheep grazing and cutting (Research Note) 412
Wilson, F.E.A., Harvey, B.M.R., McAdam, J.H. and Walton, D.W.H. The response of Whitegrass (*Cortaderia pilosa* (D'Urv.) Hack.) to nitrogen nutrition 84
Winters, A. *see* Fraser, M.D.
Winters, A.L., Fychan, R. and Jones, R. Effect of formic acid and a bacterial inoculant on the amino acid composition of grass silage and on animal performance 181
Wright, I.A., Jones, J.R. and Parsons, A.J. Effects of grazing by sheep or cattle on sward structure and subsequent performance of weaned lambs 138

Yarrow, N.H. and Penning, P.D. The liveweight gain of Limousin x Friesian heifers grazing perennial ryegrass/white clover swards of different clover content and the effects of their grazing on sward botanical composition 238
Youell, S.J. *see* Dewhurst, R.J.

Zenoff, A. *see* Córdoba, A.